

**KEYSTONE MOBILE PARTNERS
225 NEWTOWN RD, WARMINSTER, PA 18974
215-675-9900 – FAX 215-675-4096**

Extracorporeal Shock Wave Lithotripsy Patients:

Please follow these special instructions concerning Aspirin, Ibuprofen, Prescription Blood Thinners, Vitamins and Herbal Supplements.

If you have any questions regarding any medications, please contact your physician, pharmacists or please feel free to call us directly and ask to speak with a nurse at 215-675-9900 M-F 9-5.

!!ASPIRIN (81-325-650MG), ASPIRIN CONTAINING PRODUCTS, MOTRIN, ADVIL, IBUPROFEN OR NSAID CONTAINING PRODUCTS MUST BE STOPPED 7 DAYS PRIOR TO TREATMENT. FAILURE TO STOP THESE MEDICATIONS WILL RESULT IN THE CANCELLATION OF YOUR PROCEDURE!!

The first stopped day is the first day that you DID NOT take the medication. The last day is the day BEFORE your treatment.

PRESCRIPTION ANTICOAGULANT MEDICATION:

The following list is the minimum required number of days to stop these medications prior to your treatment. Please check with your prescribing physician to determine if it will be safe for you to stop these medications for the number of listed days. If it is deemed unsafe, by your prescribing physician, to stop these medications, you will not be a candidate for (ESWL) lithotripsy. These medications **MUST** be stopped for the number of listed days prior to your treatment. **Failure to stop these medications will result in the cancellation of your procedure.**

Aggrenox (Dipyridamole/Aspirin) – **STOP 7 DAYS**
Brilinta (Ticagrelor) – **STOP 5 DAYS**
Coumadin (Require PT/INR \leq 1.1) – **STOP 5 DAYS**
Eliquis (Apixaban) – **STOP 3 DAYS**
Effient (Prasugrel) – **STOP 7 DAYS**
Lovenox – **STOP 24 HOURS**
Persantine (Dipyridamole) – **STOP 24 HOURS**

Plavix – **STOP 7 DAYS**
Pletal – **STOP 5 DAYS**
Pradaxa – **STOP 3 DAYS**
Savaysa (Edoxaban) – **STOP 3 DAYS**
Ticlid (Ticlopidine HCL) – **STOP 10 DAYS**
Xarelto – **STOP 2 DAYS**

The following list of medications contain ingredients that may interfere with your blood's ability to clot. **You must discontinue the use of these medications 7 days prior to your lithotripsy procedure.**

!! FAILURE TO STOP THESE MEDICATIONS WILL RESULT IN THE CANCELLATION OF YOUR PROCEDURE!!

Please be advised that ALL medications should be discussed with your physician prior to lithotripsy. These include ALL prescription medications and ALL over-the-counter (OTC) medications.

NONSTEROIDAL ANTIINFLAMMATORY MEDICATIONS (NSAID):

There are many over-the-counter (OTC) products that contain this type of medicine and if your medication is not listed please consult your physician as to the need to stopping prior to your treatment

GENERIC NAME	BRAND NAME
Celecoxib	Celebrex
Diclofenac	Arthotec, Cambia, Cataflam, Dyloject, Flector, Pennsaid, Solaraze, Volataren, Voltaren XR, Zipsor, Zorvolex,
Diflunisal	Dolobid
Etodolac	Lodine, Lodine XL
Fenoprofen	Fenortho, Nalfon, Nalfon 200, ProFen
Flurbiprofen	Ansaid
Ibuprofen	Actiprofem, Addaprin, Advil (all varieties), Caldolor, Combunox, Dristan Sinus, Duexis, Genpril, Haltran, IBU (all varieties), Ibuprohm, I-Prin, Medipren, Menadol, Midol IB, Motrin (all varieties), NeoProfenm, Nuprin, Proprinal, Q-Profen, Reprexain, Rufen, Saletto (all varieties), Trendar, Vicoprofen
Indomethacin	Indocin, Indocin SR, Indo-Lemmon, Indomethagan, Tivorbex
Ketorolac	Sprix, Toradol
Ketoprofen	Orudis, Orudis KT, Oruvail
Mefenamic Acid	Meclodium, Meclofenamate, Meclomen, Ponstel
Meloxicam	Mobic, Omiiz ODT, Vivlodex
Nabumetone	Ralafen
Naproxen	Aflaxen, Aleve (all varieties), All Day Pain Relief, Anaprox, Anaprox DS, Flanax Pain Reliever, Midol ER, Naprelan, Naprosyn, Naprosyn EC, Naproxen, Prevacid Naprapac, Treximet, Vimovo
Oxaprozin	Daypro
Piroxicam	Feldene
Sulindac	Clinoril
Tolmetin	Tolectin, Tolectin 600, Tolectin DS

The following list of medications contain ingredients that may interfere with your blood's ability to clot. **You must discontinue the use of these medications 7 days prior to your lithotripsy procedure.**

!! FAILURE TO STOP THESE MEDICATIONS WILL RESULT IN THE CANCELLATION OF YOUR PROCEDURE!!

Please be advised that ALL medications should be discussed with your physician prior to lithotripsy. These include ALL prescription medications and ALL over-the-counter (OTC) medications.

ASPIRIN (ASA) and SALICYLATE MEDICATIONS:

*There are many medications that contain Aspirin (ASA) and Salicylate under many different names. If your medication is not listed, please consult your physician as to the need to stopping prior to your treatment. If you are not sure if your medication contains Aspirin or Salicylate please contact your physician or pharmacists to determine if your medication contains either of those ingredients. *

ASPIRIN (ASA) or ASPRIN COMPOUND MEDICATIONS	ASPIRIN (ASA) or ASPRIN COMPOUND MEDICATIONS	MAGNESIUM SALICYLATE
Acetylsalicylic Acid	Lortab ASA	Doan's Pills (all varieties)
Acuprin	Meprobamate Aspirin (Equagesic)	Choline Magnesium Trisalicylate
Aggrenox	Norgesic (all varieties)	Mobidin
Alka Seltzer (all varieties)	Orphenadrine	Mobigesic
Anacin (all varieties)	Orphengesic	Nuprin Backache
Arthritis Pain	Pamprin (all varieties)	Tricosal
Ascriptin (all varieties)	Percodan	Trilisate
Asper-gum	Roxiprin	
Aspirin (all varieties, all brands (name or store brand), ALL doses, enteric or non-enteric coated	Soma Compound	SALSALATE
Aspirin/Oxycodone (all brands)	St. Joseph Aspirin (all varieties)	Disalcid
Asprimox	Synalgos-DC	Salflex
Bayer Aspirin (all varieties)	Trigesic	Salsitab
Bufferin (all varieties)	Vanquish	Sodium Salicylate
Darvon-N-Compound	Zorprin	
Doxaphene Compound		
Durlaza		
Ecotrin (all varieties)		
Empirin/Codeine		
Entercote		
Fiornial		
Genacote		

The following list of vitamins and herbal supplements contain ingredients that may interfere with your blood's ability to clot. **You must discontinue the use of these medications/supplements 7 days prior to your lithotripsy procedure.**

Please be advised that ALL medications should be discussed with your physician prior to lithotripsy. These include ALL prescription medications and ALL over-the-counter (OTC) medications.

There are many over-the-counter (OTC) products that contain ingredients that can cause bleeding during and after surgery and if your medication/supplement is not listed please consult your physician as to the need to stopping prior to your treatment

VITAMINS and HERBAL SUPPLEMENTS:

Dehydroepiandrosterone (DHEA)

Eicosapentaenoic Acid (EPA)

Fish Oils (Omega-3 Fatty Acids)

Garlic Supplement

Lovaza

Onion Supplement

Vitamin E